arXiv:1305.0071v1 [hep-th] 1 May 2013

Cohomology ring of the BRST operator associated to the sum of two pure spinors

A. Mikhailov[a], A. Schwarz[b], Renjun Xu[c,*]

[a] *Instituto de Física Teórica, Universidade Estadual Paulista*
R. Dr. Bento Teobaldo Ferraz 271, Bloco II – Barra Funda
CEP:01140-070 – São Paulo, Brasil
[b] *Department of Mathematics, University of California, Davis, CA 95616, USA*
[c] *Department of Physics, University of California, Davis, CA 95616, USA*

Abstract

In the study of the Type II superstring, it is useful to consider the BRST complex associated to the sum of two pure spinors. The cohomology of this complex is an infinite-dimensional vector space. It is also a finite-dimensional algebra over the algebra of functions of a single pure spinor. In this paper we study the multiplicative structure.

The central object of the pure spinor formalism is the BRST operator Q, which involves nonlinearly constrained ghosts. For example, the Type IIB superstring uses two ghosts λ_L and λ_R, which are spinors of $so(10)$ satisfying the *pure spinor constraint*:

$$(\lambda_L \Gamma^m \lambda_L) = (\lambda_R \Gamma^m \lambda_R) = 0 \tag{1}$$

Roughly speaking, the BRST structures are in one-to-one correspondence with SUGRA backgrounds. For a given background, the cohomology of Q describes its infinitesimal deformations. It was shown in [1, 2] that the knowledge of the cohomology of the simplified differential $Q^{(0)} = (\lambda_L + \lambda_R)\frac{\partial}{\partial\theta}$ where θ is an odd spinor variable is useful for understanding the cohomology of Q.

The cohomology groups of $Q^{(0)}$ were calculated in [2] as modules over the ring of polynomials of $\lambda_L^\alpha, \lambda_R^\alpha$. In the present paper we will study the multiplication of the cohomology classes, and calculate the cohomology as a ring. This allows us to simplify the description of cohomology. We can use the multiplication in cohomology to obtain information about cohomology of more complicated differentials.

Let us consider the ring $A = \mathbb{C}[\lambda_L, \lambda_R, \theta]$ of polynomials depending of ten-dimensional even pure spinors λ_L, λ_R and a ten-dimensional odd spinor θ. In other words, this means that the components λ_L, λ_R obey the pure spinor constraint (1) and the components θ^α are free Grassmann variables.

We define a differential acting on A by the formula:

$$Q = (\lambda_L^\alpha + \lambda_R^\alpha)\,\frac{\partial}{\partial\theta^\alpha} \tag{2}$$

*Corresponding author.
Email address: rxu@ucdavis.edu (Renjun Xu)

This differential commutes with the multiplication by a polynomial of λ_L, λ_R. Let R denote the ring of polynomials of λ_L and λ_R:

$$R = \mathbf{C}[\lambda_L, \lambda_R] \tag{3}$$

Then A is a DGA over R, *i.e.* a differential graded R-algebra. The grading is introduced as the number of thetas (we will use the term θ-degree for this grading). As Q commutes with the elements of R, the cohomology $H(A)$ can also be regarded as a graded R-algebra. The natural action of $so(10)$ on A induces a representation of $so(10)$ on $H(A)$. The permutation of λ_L and λ_R commutes with the differential and therefore induces an involution on $H(A)$. Our goal is to calculate $H(A)$ with all of these structures.

First of all we describe the elements that generate $H(A)$ as a unital R-algebra. This means that all cohomology classes can be obtained from them and the unit element by means of multiplication, addition and multiplication by a polynomial of λ_L, λ_R.

We prove that $H(A)$ is generated by the cohomology classes of the following elements:

$$\Phi^p = (\lambda_L - \lambda_R)\Gamma^p\theta \ , \quad \deg \Phi^p = 1, \tag{4}$$

$$\Phi \ = (\lambda_L\Gamma^p\theta)(\lambda_R\Gamma^p\theta) \ , \quad \deg \Phi = 2, \tag{5}$$

$$\Psi_\alpha = \frac{\partial}{\partial\theta^\alpha}\Psi \ , \quad \deg \Psi_\alpha = 4, \tag{6}$$

$$\Psi \ = \left(1 - \frac{5}{3}\left(\lambda_L\frac{\partial}{\partial\lambda_R}\right)\right)(\lambda_R\Gamma^p\theta)(\lambda_R\Gamma^q\theta)(\lambda_R\Gamma^r\theta)(\theta\Gamma_{pqr}\theta) \ - \ \{R \to L\} \ , \tag{7}$$
$$\deg \Psi = 5$$

Here $\frac{\partial}{\partial\lambda_R}$ stands for a formal differentiation with respect to λ_R, without taking into account the pure spinor constraint. The second term in (7) is obtained from the first one by interchanging L and R.

Using the algebra generators we can write down elements that generate $H(A)$ as an R-module. Namely, we should add to them the unit element of the algebra, 45 elements Φ^{ab} and 120 elements Φ^{abc} given by formulas

$$\Phi^{ab} = \Phi^a\Phi^b \tag{8}$$

$$\Phi^{abc} = \Phi^a\Phi^b\Phi^c \tag{9}$$

In other words, all elements of $H(A)$ can be obtained as linear combinations of $1, \Phi^p, \Phi$, Ψ_α, Ψ, Φ^{ab}, Φ^{abc} with coefficients belonging to $R = \mathbb{C}[\lambda_L, \lambda_R]$. The $so(10)$-action on $H(A)$ is determined by the action on ring generators. The generators Φ and Ψ are scalars; ten generators Φ^a as a vector $V = [1, 0, 0, 0, 0]$; sixteen generators Ψ_α transform according the dual spinor representation $S^* = [0, 0, 0, 1, 0]$. It follows that Φ^{ab} transforms as $\Lambda^2 V = [0, 1, 0, 0, 0]$ and Φ^{abc} as $\Lambda^3 V = [0, 0, 1, 0, 0]$

We can consider $H(A)$ as a vector space, or as a representation of $so(10)$. We can introduce an additional grading as the number of lambdas. We denote:

- $H^{N,q}(A)$ the component of $H(A)$ with λ-grading $(N - q)$ and θ-grading q. (Hence N stands for total number of lambdas and thetas.)
- $H^q(A)$ the component of $H(A)$ consisting of elements of θ-degree q (*i.e.* the sum of $H^{N,q}(A)$ over all possible N).

We can analyze the structure of $H^{N,q}(A)$ as $so(10)$-representation. We obtain

$$\begin{aligned} H^{N,0} &= [0,0,0,0,N] && (10)\\ H^{N,1} &= [1,0,0,0,N-2] && (11)\\ H^{N,2} &= [0,1,0,0,N-4], \text{ when } N \neq 4 && (12)\\ H^{N,3} &= [0,0,1,0,N-6] && (13)\\ H^{N,4} &= [0,0,0,1,N-7] && (14)\\ H^{N,5} &= [0,0,0,0,N-8] && (15) \end{aligned}$$

When $N = 4$, there is one additional term, a scalar, in $H^{4,2}$:

$$H^{4,2} = [0,0,0,0,0] \oplus [0,1,0,0,0] \tag{16}$$

Notice that the involution of $H(A)$ corresponding to the interchange of λ_L and λ_R acts as a multiplication by $(-1)^{N-q}$ on almost all elements of $H^{N,q}$, the only exception being $[0,0,0,0,0]$ in $H^{4,2}$ where it acts as the multiplication by -1. Sometimes it is more convenient to use the involution $\lambda_L \to -\lambda_R$, $\lambda_R \to -\lambda_L$ combining the interchange of lambdas and the sign change; the groups $H^{N,q}$ with the exception of $[0,0,0,0,0]$ in $H^{4,2}$ are invariant with respect to this involution.

The vector space spanned by the generators of the R-module to $H^q(A)$ will be denoted by W^q; this space carries a representation of $so(10)$. This representation is irreducible in all cases except for $q = 2$. It is easy to check that:

$$\begin{aligned} W^1 = H^{2,1} &= [1,0,0,0,0],\\ W^2 = H^{4,2} = [0,0,0,0,0] \oplus [0,1,0,0,0] &= W_1^2 + W_2^2,\\ W^3 = H^{6,3} &= [0,0,1,0,0],\\ W^4 = H^{7,4} &= [0,0,0,1,0],\\ W^5 = H^{8,5} &= [0,0,0,0,0]. \end{aligned}$$

Some elements of R act as zero on $H(A)$. Namely, consider the ideal of R consisting of the polynomials of the form $(\lambda_L^\alpha + \lambda_R^\alpha)P_\alpha(\lambda_L, \lambda_R)$ where $P_\alpha(\lambda_L, \lambda_R)$ can be any polynomial. This ideal acts as zero on $H(A)$, because:

$$(\lambda_L^\alpha + \lambda_R^\alpha)P_\alpha(\lambda_L, \lambda_R) = Q\left(\theta^\alpha P_\alpha(\lambda_L, \lambda_R)\right) \tag{17}$$

Let us denote R_+ the factorspace of R over this ideal. The degree k component R_+^k carries the representation $[0,0,0,0,k]$ of $so(10)$. Geometrically, we can interpret R_+ as the algebra of functions of a single pure spinor. We see that $H^q(A)$ is actually an R_+-module. In other words, we have a map

$$a \ : \ R_+^k \otimes W^q \to H^{k+q+l(q),q} \tag{18}$$

where $l(q)$ stands for the λ-degree of W^q. One can check that this map is surjective (moreover, for $q = 2$ this statement remains correct if $W^{q=2}$ is replaced with W_2^2; in fact $R_+^{k>0} \otimes W_1^2 \subset \ker a$). In other words, all cohomology classes can be obtained as linear combinations of generators with coefficients from R_+.

Using the above results we can find the bases of vector spaces $H^{N,q}(A)$. For example, one can find the highest weight vectors of the representations acting on these spaces; then one can construct

a basis applying the operators corresponding to negative roots. The highest weight vector of $H^{N,q}$ can be obtained as a product of the highest weight vector of the k-th graded component of the ring R_+ and the highest weight vector of W^q. (Here $k = N - q - l(q)$.) For example, when $q = 2$, the highest weight vector of $H^{N,2}$ can be obtained as a product of $(\lambda_L^1 - \lambda_R^1)^{N-4}$ and the vector $\Phi^1\Phi^2$ in W^2. [1]

Our results can be applied to the analysis of the cohomology of the differential Q_{ch} that is obtained from Q adding the expression $((\lambda_L - \lambda_R)\Gamma^p\theta)\frac{\partial}{\partial x^p}$. (This operator acts in the space of polynomial functions of $\lambda_L, \lambda_R, \theta$ and x. One can consider instead of polynomials of x other classes of functions.) Using the notations above we can represent Q_{ch} in the form

$$Q_{ch} = Q + \Phi^p \frac{\partial}{\partial x^p}.$$

To calculate the cohomology of Q_{ch} we can apply the spectral sequence of bicomplex (in more physical words, we can consider the second summand as a perturbation). The first approximation (the E_2-term of spectral sequence) is given by the cohomology of the differential induced by the second summand on the cohomology of the first summand (on $H(A) \otimes \mathbb{C}[x]$). To compute this differential we should calculate the product of Φ^p with generators of modules $H^q(A)$. We obtain

$$\Phi^p\Phi^q = \Phi^{pq} \tag{19}$$

$$\Phi^p\Phi = 0 \tag{20}$$

$$\Phi^p\Phi^{ql} = \Phi^{pql} \tag{21}$$

$$\Phi^p\Phi^{qlm} \simeq (\Gamma^{pqlm})^\alpha_\beta(\lambda_L^\beta - \lambda_R^\beta)\Psi_\alpha \tag{22}$$

$$\Phi^p\Psi_\alpha \simeq \Gamma^p_{\alpha\beta}(\lambda_L^\beta - \lambda_R^\beta)\Psi \tag{23}$$

$$\Phi^p\Psi = 0 \tag{24}$$

We have not calculated the coefficients in (22) and (23).

We can calculate also the complete multiplication table of generators of $H(A)$ considered as R-module.

Let us describe the way we obtained our results. Using the Macaulay2 package [3] we can calculate the number of generators of the R-algebra $H(A)$, the degrees of these generators, and the number of generators of the R-modules $H^k(A)$. We find that the R-algebra $H(A)$ is generated by 28 elements (ten of degree 1, one of degree 2, sixteen of degree 4, and one of degree 5), the number of generators of $H^q(A)$ is equal to 195 (with one of degree 0, ten of degree 1, 46 of degree 2, 120 of degree 3, sixteen of degree 4, and one of degree 5). The calculation gives us also the Hilbert series of $H(A)$ (the generating function of $dimH^{N,q}$); we obtain

$$\sum dimH^{N,q}\tau_0^N\tau_1^q = \sum_{q=0}^{5} P_q(\tau_0)\tau_1^q,$$

[1] Here we use a convention of choosing λ^1 and $\Phi^1\Phi^2$ as the highest weight vectors in spinor representation and in $\Lambda^2 V$ respectively

where

$$
\begin{aligned}
P_0(\tau_0) &= \frac{1+5\tau_0+5{\tau_0}^2+{\tau_0}^3}{(1-\tau_0)^{11}} \\
P_1(\tau_0) &= \frac{10{\tau_0}^2+34{\tau_0}^3+16{\tau_0}^4}{(1-\tau_0)^{11}} \\
P_2(\tau_0) &= (46{\tau_0}^4+54{\tau_0}^5+66{\tau_0}^6-166{\tau_0}^7+330{\tau_0}^8-462{\tau_0}^9+462{\tau_0}^{10}-330{\tau_0}^{11} \\
&\quad +165{\tau_0}^{12}-55{\tau_0}^{13}+11{\tau_0}^{14}-{\tau_0}^{15})/(1-\tau_0)^{11} \\
P_3(\tau_0) &= (120{\tau_0}^6-120{\tau_0}^7+330{\tau_0}^8-462{\tau_0}^9+462{\tau_0}^{10}-330{\tau_0}^{11}+165{\tau_0}^{12} \\
&\quad -55{\tau_0}^{13}+11{\tau_0}^{14}-{\tau_0}^{15})/(1-\tau_0)^{11} \\
P_4(\tau_0) &= \frac{16{\tau_0}^7+34{\tau_0}^8+10{\tau_0}^9}{(1-\tau_0)^{11}} \\
P_5(\tau_0) &= \frac{{\tau_0}^8+5{\tau_0}^9+5{\tau_0}^{10}+{\tau_0}^{11}}{(1-\tau_0)^{11}}
\end{aligned}
$$

To find the decomposition of $H^{N,q}$ into direct sum of irreducible representations of $so(10)$ we use the Lie code [4] that allows to find such a decomposition for the graded components $A^{N,q}$ of A for small N, q. Applying Schur's lemma and the information about the Hilbert series we can find the action of the differential on the irreducible components of $A^{N,q}$; this allows us to find the $so(10)$-action on $H^{N,q}$ for small N, q. (We use a version of the "maximal propagation principle"[5],[6] in this consideration.) These results allow us to guess the representatives of the cohomology classes of generators. We can check our guess using [3].

Knowing the $so(10)$-action on $H^{N,q}$ for small N, q we guess the $so(10)$ representation for all N, q; we check our guess using the formula for the dimension of representation and the Hilbert series. To give a rigorous proof we can verify that the representation we guessed is embedded in $H^{N,q}$.

Acknowledgements We are indebted to N. Berkovits and M. Movshev for interesting discussions.

References

[1] N. Berkovits and Th. Fleury, *Harmonic Superspace from the $AdS_5 \times S^5$ Pure Spinor Formalism*, *JHEP* **1303** (2013) 022, [arXiv/1212.3296]

[2] A. Mikhailov and R. Xu, *BRST cohomology of the sum of two pure spinors*, [arXiv/1301.3353].

[3] D. R. Grayson and M. E. Stillman, "Macaulay 2, a software system for research in algebraic geometry." http://www.math.uiuc.edu/Macaulay2/.

[4] A. M. Cohen, B. Lisser, and M. A. A. v. Leeuwen, "LiE: A Computer algebra package for Lie group computations." http://www-math.univ-poitiers.fr/~maavl/LiE/.

[5] M. Movshev, A. Schwarz, and R. Xu, *Homology of Lie algebra of supersymmetries and of super Poincare Lie algebra*, *Nuclear Physics B* **854** (2011) 483–503 doi: **10.1016/j.nuclphysb.2011.08.023** [arXiv/1106.0335].

[6] M. Cederwall, B. E. Nilsson, and D. Tsimpis, *Spinorial cohomology and maximally supersymmetric theories*, *JHEP* **0202** (2002) 009 [arXiv/hep-th/0110069].